

Common Name	Scientific Name	Disjunct	History	BC Status	COSEWIC	Region
TERRESTRIAL MAMMALS						
Arctic Shrew	<i>Sorex arcticus (arcticus)</i>			Yellow		Northeast, east of the Rocky Mountains from the Peace River to Kotcho Lake.
Pacific Water Shrew ^S	<i>Sorex bendirii</i>			Red	T	Lower Fraser River Valley, west of Chilliwack River and Agassiz, south of the north shore of Burrard Inlet.
Merriam's Shrew	<i>Sorex merriami</i>		RR	Red		South Okanagan.
Preble's Shrew ^S	<i>Sorex preblei</i>		RR	Red		South Okanagan.
Trowbridge's Shrew ^S	<i>Sorex trowbridgii (trowbridgii)</i>			Blue		Lower Fraser River valley.
Tundra Shrew ^S	<i>Sorex tundrensis (tundrensis)</i>		SR	Red		Chilkat Pass and Kelsall Lake on the Haines Road.
American Shrew Mole ^S	<i>Neurotrichus gibbsii (gibbsii)</i>			Yellow		Fraser River Delta and Cascade Mountains east to Manning Provincial Park, north to Sechelt Peninsula and Pitt Lake.
Coast Mole ^S	<i>Scapanus orarius (schefferi)</i>			Yellow		Lower Fraser River basin, east to Hope and north to south shore of Burrard Inlet.
Townsend's Mole ^S	<i>Scapanus townsendii (townsendii)</i>			Red	E	Near Huntingdon and Abbotsford adjacent to the US border.
Spotted Bat	<i>Euderma maculatum</i>			Blue	SC	Low elevations in interior valleys.
Western Red Bat	<i>Lasiurus blossevillei (frantzi)</i>	D1	SR	Red		Skagit Valley and Okanagan Valley.

Stewardship responsibility at the species level is given below subspecies scientific name if species is of intermediate (11%-29% range or population in BC) or high (>30% range or population in BC) stewardship responsibility. Otherwise, all species are wide-ranging with <10% global range or population over >30% of BC.

Common Name	Scientific Name	Disjunct	History	BC Status	COSEWIC	Region
TERRESTRIAL MAMMALS						
Pallid Bat	<i>Antrozous pallidus (pallidus)</i>			Red	T	South Okanagan Valley, between Okanagan Falls and Osoyoos.
Townsend's Big-eared Bat	<i>Corynorhinus townsendii</i>			Blue		Vancouver Island and adjacent islands, Southern Interior north to Williams Lake, east to Creston.
	<i>Corynorhinus townsendii pallescens</i>			NR (Blue)		Southern Interior.
	<i>Corynorhinus townsendii townsendii</i>			NR (Blue)		Southwest coast.
California Myotis	<i>Myotis californicus californicus</i>			NR (Yellow)		Southern Interior from Okanagan Valley to Revelstoke.
Western Small-footed Myotis	<i>Myotis ciliolabrum (melanorhinus)</i>			Blue		Okanagan and Similkameen Valleys.
Little Brown Myotis	<i>Myotis lucifugus carissima</i>			NR (Yellow)		Southern Interior, Okanagan Valley to Kamloops.
	<i>Myotis lucifugus lucifugus</i>			NR (Yellow)		Extreme north.
Northern Long-eared Myotis	<i>Myotis septentrionalis</i>			Blue		A few locations in east.
Yuma Myotis	<i>Myotis yumanensis sociabilis</i>			NR (Yellow)		Southern Interior north to Kamloops, east to Creston.
Collared Pika ^S	<i>Ochotona collaris</i>			Yellow		Extreme northwest.
American Pika ^S	<i>Ochotona princeps brunnescens</i> Species intermediate			NR (Yellow)		Coastal mountains, north to Garibaldi Provincial Park.

Stewardship responsibility at the species level is given below subspecies scientific name if species is of intermediate (11%-29% range or population in BC) or high (>30% range or population in BC) stewardship responsibility. Otherwise, all species are wide-ranging with <10% global range or population over >30% of BC.

Common Name	Scientific Name	Disjunct	History	BC Status	COSEWIC	Region
TERRESTRIAL MAMMALS						
Snowshoe Hare	<i>Lepus americanus bairdii</i>			NR (Yellow)		Extreme southeast.
	<i>Lepus americanus cascadenis</i>			NR (Yellow)		Coastal, north to Jervis Inlet.
	<i>Lepus americanus dalli</i>			NR (Yellow)		North.
	<i>Lepus americanus washingtonii</i>			Red		Lower Fraser Valley.
Mountain Cottontail	<i>Sylvilagus nuttallii (nuttallii)</i>			Blue	(SC)	South Okanagan, Osoyoos to Summerland and in the Similkameen Valley west of Keremeos.
Mountain Beaver ^S	<i>Aplodontia rufa</i>					Lower Fraser Valley and Cascades to Nicola Valey.
	<i>Aplodontia rufa rufa</i>			Red	SC	Lower Fraser Valley.
Southern Red-backed Vole	<i>Clethrionomys gapperi athabascaae</i>			NR (Yellow)		Northeast.
	<i>Clethrionomys gapperi galei</i>			Blue		Southeast, Sage and Kishinena drainages.
	<i>Clethrionomys gapperi occidentalis</i>	SR, H		Red		Burns Bog.
Northern Red-backed Vole	<i>Clethrionomys rutilus (dawsonii)</i>			Yellow		Extreme northwest, Haines Triangle to Summit Pass.
Long-tailed Vole	<i>Microtus longicaudus littoralis</i>			NR (Yellow)		Stikine River area.
	<i>Microtus longicaudus longicaudus</i>			NR (Yellow)		Southern dry interior, north to Kamloops and Shuswap.
Montane Vole	<i>Microtus montanus</i>			Yellow		Southern Interior, Okanagan Valley to Williams Lake.
Tundra Vole	<i>Microtus oeconomus</i>			Yellow		Extreme northwest BC, Haines Triangle and Cassiar Mountains.

Stewardship responsibility at the species level is given below subspecies scientific name if species is of intermediate (11%-29% range or population in BC) or high (>30% range or population in BC) stewardship responsibility. Otherwise, all species are wide-ranging with <10% global range or population over >30% of BC.

Common Name	Scientific Name	Disjunct	History	BC Status	COSEWIC	Region
TERRESTRIAL MAMMALS						
Creeping Vole ^S	<i>Microtus oregoni</i>			Yellow		Vancouver to Allison Pass, Manning Provincial Park.
Meadow Vole	<i>Microtus pennsylvanicus modestus</i>			NR (Yellow)		South, southern Okanagan to Rocky Mountain Trench.
Townsend's Vole ^S	<i>Microtus townsendii townsendii</i> Species 30-50% range in BC			NR (Yellow)		Southwest, north to Burrard Inlet, east to Chilliwack.
Eastern Heather Vole	<i>Phenacomys ungava</i>			NR		North.
Northern Bog Lemming	<i>Synaptomys borealis borealis</i>			NR (Yellow)		Northeast.
Beaver	<i>Castor canadensis belugae</i>			NR (Yellow)		Coastal, from Alaska border to Dean Inlet.
	<i>Castor canadensis canadensis</i>			NR (Yellow)		Northeast.
Bushy-tailed Woodrat	<i>Neotoma cinerea cinerea</i> Species 30-50% range in BC			NR (Yellow)		Extreme southeast.
Deer Mouse	<i>Peromyscus maniculatus artemisiae</i>			NR (Yellow)		South central, north to Prince George.
Western Harvest Mouse	<i>Reithrodontomys megalotis (megalotis)</i>			Blue	(SC)	Grassland in the Okanagan and lower Similkameen Valleys.
Porcupine	<i>Erethizon dorsatum myops</i>			NR (Yellow)		Extreme north and northeast.
Northern Pocket Gopher	<i>Thomomys talpoides</i>			Yellow		Southern Interior, southern Kootenays and southern Rocky Mountains.
	<i>Thomomys talpoides fuscus</i>			NR (Yellow)		Southern Monashee Mountains.
	<i>Thomomys talpoides saturatus</i>			NR (Yellow)		Southern Kootenays.
Great Basin Pocket Mouse ^S	<i>Perognathus parvus</i>			Blue		South and north Okanagan Valley, lower Similkameen Valley, Thompson and Kettle Valleys.

Stewardship responsibility at the species level is given below subspecies scientific name if species is of intermediate (11%-29% range or population in BC) or high (>30% range or population in BC) stewardship responsibility. Otherwise, all species are wide-ranging with <10% global range or population over >30% of BC.

Common Name	Scientific Name	Disjunct	History	BC Status	COSEWIC	Region
TERRESTRIAL MAMMALS						
Northern Flying Squirrel	<i>Glaucomys sabrinus fuliginosus</i>			NR (Yellow)		South Coast Mountains.
	<i>Glaucomys sabrinus oregonensis</i>			NR (Yellow)		Coastal, Loughborough Inlet south to US border.
Hoary Marmot	<i>Marmota caligata caligata</i> Species 30-50% range in BC			NR (Yellow)		Extreme northwest, Haines Triangle to Stikine River.
Yellow-bellied Marmot	<i>Marmota flaviventris (avara)</i>			Yellow		Southern Interior north to William's Lake.
Woodchuck	<i>Marmota monax</i>			Yellow		Extreme northeast and northcentral, Peace River area, Stikine River southeasterly to Creston.
	<i>Marmota monax canadensis</i>			NR		Peace River region.
	<i>Marmota monax ochracea</i>			NR		Extreme north-central (Liard River) and northwest (Haines Triangle).
Golden-mantled Ground Squirrel	<i>Spermophilus lateralis</i>			Yellow		Selkirk, Purcell, Cariboo, and Rocky Mountains, north to Mount Selwyn.
Arctic Ground Squirrel	<i>Spermophilus parryii</i>			Yellow		Northwest, south to the Spatsizi and Klappan Rivers.
Douglas' Squirrel ^S	<i>Tamiasciurus douglasii mollipilosus</i> Species neither			NR - Yellow		Coastal, north to Rivers Inlet.
Red Squirrel	<i>Tamiasciurus hudsonicus petulans</i>			NR (Yellow)		Extreme northwest, Haines Triangle area.
	<i>Tamiasciurus hudsonicus preblei</i>			NR (Yellow)		Peace River-Fort Nelson region.
	<i>Tamiasciurus hudsonicus richardsoni</i>			NR (Yellow)		Rocky, Purcell and Selkirk mountains, north to Field.

Stewardship responsibility at the species level is given below subspecies scientific name if species is of intermediate (11%-29% range or population in BC) or high (>30% range or population in BC) stewardship responsibility. Otherwise, all species are wide-ranging with <10% global range or population over >30% of BC.

Common Name	Scientific Name	Disjunct	History	BC Status	COSEWIC	Region
TERRESTRIAL MAMMALS						
Least Chipmunk	<i>Neotamias minimus</i>			Yellow		North and extreme southern Rocky Mountains.
	<i>Neotamias minimus borealis</i>			NR		Northeast and north central, north to Babine Lake and Peace River.
Red-tailed Chipmunk ^S	<i>Neotamias ruficaudus ruficaudus</i>			Red		Extreme southeast; Akamina-Kishinena Provincial Park and Kootenay Pass.
Townsend's Chipmunk ^S	<i>Neotamias townsendii</i>			Yellow		Extreme southwest.
	<i>Neotamias townsendii cooperi</i>			NR		Cascades.
	<i>Neotamias townsendii townsendii</i>			NR		Lower Fraser Valley.
Meadow Jumping Mouse	<i>Zapus hudsonius alascensis</i>			Blue		Extreme northwest, Haines triangle area.
Western Jumping Mouse	<i>Zapus princeps idahoensis</i>			NR (Yellow)		Extreme southeast, Rocky Mountains and Rocky Mountain Trench north to Mount Assiniboine.
Pacific Jumping Mouse ^S	<i>Zapus trinotatus (trinotatus)</i>			Yellow		Extreme southwest, except Vancouver Island, north to Garibaldi Provincial Park, east to Allison Pass in Manning Provincial Park.
Red Fox	<i>Vulpes vulpes macroura</i>			NR (Yellow)		Southeast.
Bobcat	<i>Lynx rufus fasciatus</i>			NR (Yellow)		Coast to Bute Inlet.
Marten	<i>Martes americana arctuosa</i>			NR (Yellow)		North.
Striped Skunk	<i>Mephitis mephitis spissigrada</i>			NR (Yellow)		Lower Fraser Valley.
Long-tailed Weasel	<i>Mustela frenata nevadensis</i>			NR (Yellow)		South-central BC from coastal mountains to Kootenay Lake.

Stewardship responsibility at the species level is given below subspecies scientific name if species is of intermediate (11%-29% range or population in BC) or high (>30% range or population in BC) stewardship responsibility. Otherwise, all species are wide-ranging with <10% global range or population over >30% of BC.

Common Name	Scientific Name	Disjunct	History	BC Status	COSEWIC	Region
TERRESTRIAL MAMMALS						
Western Spotted Skunk	<i>Spilogale gracilis</i>			Yellow		Southwest from Garibaldi Provincial Park and the Sechelt Peninsula south to border along coast.
Raccoon	<i>Procyon lotor</i>			Yellow		Southwest and the Haida Gwaii.
Thinhorn Sheep	Ovis dalli dalli Species 30-50% range in BC			Blue		Extreme northwest
Moose	<i>Alces alces gigas</i>			NR (Yellow)		Extreme northwest.
	<i>Alces alces shirasi</i>			NR (Yellow)		Extreme southeast.
Mule Deer	<i>Odocoileus hemionus columbianus</i>			NR (Yellow)		Southwest, including Vancouver Island, along coast on west slopes of the Coast Mountains from US border to Rivers Inlet.
White-tailed Deer	<i>Odocoileus virginianus dacotensis</i>			NR (Yellow)		Peace River region.
Elk	<i>Cervus canadensis nelsoni</i> Subspecies dubious			NR (Yellow)		Southern Kootenays to the northeast, transplanted to various regions and to the Haida Gwaii.
Caribou	<i>Rangifer tarandus, Boreal population</i>			Blue	T	Northeast boreal plains.

Stewardship responsibility at the species level is given below subspecies scientific name if species is of intermediate (11%-29% range or population in BC) or high (>30% range or population in BC) stewardship responsibility. Otherwise, all species are wide-ranging with <10% global range or population over >30% of BC.

Common Name	Scientific Name	Disjunct	History	BC Status	COSEWIC	Region
MARINE MAMMALS						
Northern Fur-Seal	<i>Callorhinus ursinus</i>			Blue	NAR	Coastal, mostly migrating individuals from breeding rookeries in the Pribilof Islands Alaska to California, some over-winter.
Risso's Dolphin	<i>Grampus griseus</i>			Yellow	NAR	Uncommon.
Pygmy Sperm Whale	<i>Kogia breviceps</i>			Yellow	NAR	
Dwarf Sperm Whale	<i>Kogia sima</i>			Yellow	DD	One stranding, Bamfield, Vancouver Island.
Northern Right-Whale Dolphin	<i>Lissodelphis borealis</i>			Yellow	NAR	Offshore.
Hubb's Beaked Whale	<i>Mesoplodon carlhubbsi</i>			Blue	NAR	Strandings near Prince Rupert.
Harbour Porpoise	<i>Phocoena phocoena</i>			Blue	SC	Coastal.
Cuvier's Beaked Whale	<i>Ziphius cavirostris</i>			Yellow	NAR	Strandings.
False killer whale	<i>Pseudorca crassidens</i>		SR	Accidental	NAR	First recognized in BC in 1989.
Short-beaked Common Dolphin	<i>Delphinus delphis</i>		SR	NA	NA	Single stranding in 1953.
Long-beaked Common Dolphin	<i>Delphinus capensis</i>		RR	NA	NA	Series of sightings in 2002-03.
Pygmy Sperm Whale	<i>Kogia breviceps</i>		SR	Yellow	NAR	Single stranding in 2002.

Stewardship responsibility at the species level is given below subspecies scientific name if species is of intermediate (11%-29% range or population in BC) or high (>30% range or population in BC) stewardship responsibility. Otherwise, all species are wide-ranging with <10% global range or population over >30% of BC.